



ACC.14

TCT@ACC-12 | innovation in intervention

A2061

JACC April 1, 2014

Volume 63, Issue 12



Vascular Medicine

INCREASED MORTALITY IN ACUTE AORTIC DISSECTION PATIENTS WITH LOW ADHERENCE TO GUIDELINE RECOMMENDATIONS FOR IMAGING AND CLINIC FOLLOW-UP

Poster Contributions

Hall C

Saturday, March 29, 2014, 10:00 a.m.-10:45 a.m.

Session Title: Aortic and Peripheral Artery Dissections

Abstract Category: 32. Vascular Medicine: Non Coronary Arterial Disease

Presentation Number: 1105-77

Authors: *Ashish Chaddha, Kim Eagle, Himanshu Patel, G. Michael Deeb, Kevin Harris, Alan Braverman, Stuart Hutchison, Arturo Evangelista, Rossella Fattori, James Froehlich, Christoph Nienaber, Eric Isselbacher, Daniel Montgomery, Eva Kline-Rogers, Elise Woznicki, Troy LaBounty, University of Michigan, Ann Arbor, MI, USA*

Introduction: Guidelines advocate frequent follow-up after acute aortic dissection (AAD). We hypothesized that incomplete adherence may be common, and may be associated with increased mortality.

Methods: We examined individuals with AAD and follow-up for aortic imaging (n=314) and/or clinic visits (n=290). We examined rates of guideline adherence and adverse imaging findings (e.g. increased aortic size, new/expanding dissection, new aneurysm, endograft leak, enlarged false lumen, new intramural hematoma), and the relationship between follow-up and mortality.

Results: Mean age was 60.4±14.6 years, and 33.8% (106/314) were female. Type A and B AAD were noted in 46% and 54% of patients. Median follow-up 5.0 years (interquartile range 4.2-6.0), with 59 deaths. Guideline adherence for follow-up was limited, and rates of adverse imaging findings were small (Figure 1). By tertiles of follow-up (lowest to highest) for imaging, 5-year survival was 70%, 86%, and 74% (p=0.048); by tertiles of clinic visit follow-up, 5-year survival was 73%, 89%, and 72% (p=0.008). By specific intervals, improved survival was only observed in patients with imaging at 6-12 months (p=0.02), and a clinic visit within 1 month (p=0.04).

Conclusion: Imaging and clinic follow-up after AAD is frequently incomplete, and patients with less frequent follow-up have higher mortality. Early clinic follow-up and imaging during the interval with the highest rate of adverse image findings are associated with improved survival.

